Diabetes Medical Management Plan (DMMP)

This plan should be completed by the student's personal diabetes health care team, including the parents/guardian. It should be reviewed with relevant school staff and copies should be kept in a place that can be accessed easily by the school nurse, trained diabetes personnel, and other authorized personnel.

| Date of Plan: | This plan is valid for the current school year: | | | |
|-----------------------------|---|--------------|--|--|
| Student's Name: | Date of Birth: | | | |
| Date of Diabetes Diagnosis: | type 1 | type 2 Other | | |
| School: | School Phone Number: | | | |
| | Homeroom Teacher: | | | |
| | Phone: | | | |
| CONTACT INFORMATION | N | | | |
| Mother/Guardian: | | | | |
| | | | | |
| | | Cell: | | |
| Email Address: | | | | |
| Father/Guardian: | | | | |
| | | | | |
| | | Cell: | | |
| Email Address: | | | | |
| | | | | |
| | | | | |
| Telephone: | | | | |
| Email Address: | | ımber: | | |
| Other Emergency Contacts: | | | | |
| Name: | Relationship: | | | |
| Telephone: Home | | Cell: | | |

Diabetes Medical Management Plan (DMMP) - Page 2

CHECKING BLOOD GLUCOSE

| Target range of blood glucose: 70–130 mg/dL 70–180 mg/dL | | | | | | |
|---|--|--|--|--|--|--|
| Other: | | | | | | |
| Check blood glucose level: Before lunch Hours after lunch | | | | | | |
| 2 hours after a correction dose Mid-morning Before PE After PE | | | | | | |
| □ Before dismissal □ Other: □ As needed for signs/symptoms of low or high blood glucose □ As needed for signs/symptoms of illness Preferred site of testing: □ Fingertip □ Forearm □ Thigh □ Other: □ | | | | | | |
| | | | | | | Brand/Model of blood glucose meter: |
| | | | | | | Note: The fingertip should always be used to check blood glucose level if hypoglycemia is suspected. |
| | | | | | | Student's self-care blood glucose checking skills: |
| Independently checks own blood glucose | | | | | | |
| May check blood glucose with supervision | | | | | | |
| Requires school nurse or trained diabetes personnel to check blood glucose | | | | | | |
| Continuous Glucose Monitor (CGM): Yes No Brand/Model: Alarms set for: (low) and (high) | | | | | | |
| Note: Confirm CGM results with blood glucose meter check before taking action on sensor blood glucose level. If student has symptoms or signs of hypoglycemia, check fingertip blood glucose level regardless of CGM. | | | | | | |
| HYPOGLYCEMIA TREATMENT | | | | | | |
| Student's usual symptoms of hypoglycemia (list below): | | | | | | |
| | | | | | | |
| If exhibiting symptoms of hypoglycemia, OR if blood glucose level is less thanmg/dL, give a quick-acting glucose product equal to grams of carbohydrate. | | | | | | |
| Recheck blood glucose in $10-15$ minutes and repeat treatment if blood glucose level is less than $_____ mg/dL$. | | | | | | |
| Additional treatment: | | | | | | |

Diabetes Medical Management Plan (DMMP) - Page 3

HYPOGLYCEMIA TREATMENT (Continued)

| Follow physical activity and sports orders (see page 7). | | | | | | | | | | |
|--|--|--|--|--|--|--|--|--|--|--|
| If the student is unable to eat or drink, is unconscious or unresponsive, or is having seizure activity or convulsions (jerking movements), give: Glucagon: 1 mg 1/2 mg Route: SC IM Site for glucagon injection: arm thigh Other: | | | | | | | | | | |
| | | | | | | Call 911 (Emergency Medical Services) and the student's parents/guardian. Contact student's health care provider. | | | | |
| | | | | | | | | | | |
| | | | | | | | | | | |
| Check Urine Blood for ketones every hours when blood glucose levels are above mg/dL. | | | | | | | | | | |
| For blood glucose greater thanmg/dL AND at leasthours since last insulin dose, give correction dose of insulin (see orders below). | | | | | | | | | | |
| For insulin pump users: see additional information for student with insulin pump. | | | | | | | | | | |
| Give extra water and/or non-sugar-containing drinks (not fruit juices):ounces per hour. | | | | | | | | | | |
| Additional treatment for ketones: | | | | | | | | | | |
| | | | | | | | | | | |

Follow physical activity and sports orders (see page 7).

- Notify parents/guardian of onset of hyperglycemia.
- If the student has symptoms of a hyperglycemia emergency, including dry mouth, extreme thirst, nausea and vomiting, severe abdominal pain, heavy breathing or shortness of breath, chest pain, increasing sleepiness or lethargy, or depressed level of consciousness: Call 911 (Emergency Medical Services) and the student's parents/ guardian.
- Contact student's health care provider.

INSULIN THERAPY Insulin delivery device: syringe insulin pen insulin pump Type of insulin therapy at school: Adjustable Insulin Therapy Fixed Insulin Therapy ■ No insulin **Adjustable Insulin Therapy** Carbohydrate Coverage/Correction Dose: Name of insulin: Carbohydrate Coverage: Insulin-to-Carbohydrate Ratio: Lunch: 1 unit of insulin per _____ grams of carbohydrate Snack: 1 unit of insulin per grams of carbohydrate **Carbohydrate Dose Calculation Example** Grams of carbohydrate in meal = __ units of insulin Insulin-to-carbohydrate ratio • Correction Dose: Blood Glucose Correction Factor/Insulin Sensitivity Factor = _____ Target blood glucose = mg/dL**Correction Dose Calculation Example** Actual Blood Glucose-Target Blood Glucose = ____ units of insulin Blood Glucose Correction Factor/Insulin Sensitivity Factor Correction dose scale (use instead of calculation above to determine insulin correction dose): Blood glucose _____ to ____ mg/dL give ____units Blood glucose _____ to ____ mg/dL give ____ units Blood glucose _____ to ____ mg/dL give ____units Blood glucose _____ to ____ mg/dL give ____units

Diabetes Medical Management Plan (DMMP) – page 4

Diabetes Medical Management Plan (DMMP) – page 5

INSULIN THERAPY (Continued)

| When to give insu | lin: | | | |
|---|--|--|--|--|
| Lunch | | | | |
| Carbohydrate | coverage only | | | |
| Carbohydrate coverage plus correction dose when blood glucose is greater thanmg/dL and hours since last insulin dose. | | | | |
| Other: | | | | |
| Snack | | | | |
| No coverage for | or snack | | | |
| Carbohydrate | | | | |
| Carbohydrate | coverage plus correction dose when blood glucose is greater than and hours since last insulin dose. | | | |
| Other: | | | | |
| Correction dos | se only: | | | |
| | cose greater thanmg/dL AND at least hours since last | | | |
| insulin dose. | | | | |
| Other: | | | | |
| Fixed Insulin Thera | apv | | | |
| | ~P) | | | |
| _ | insulin given pre-lunch daily | | | |
| | insulin given pre-snack daily | | | |
| | misumi given pre-snack dany | | | |
| otner. | | | | |
| Parental Authoriza | ation to Adjust Insulin Dose: | | | |
| Yes No | Parents/guardian authorization should be obtained before administering a correction dose. | | | |
| Yes No | Parents/guardian are authorized to increase or decrease correction dose scale within the following range: +/ units of insulin. | | | |
| Yes No | Parents/guardian are authorized to increase or decrease insulin-to- | | | |
| les la No | carbohydrate ratio within the following range: units per prescribed grams of carbohydrate, +/ grams of carbohydrate. | | | |
| Yes No | Parents/guardian are authorized to increase or decrease fixed insulin dose within the following range: +/ units of insulin. | | | |

Diabetes Medical Management Plan (DMMP) – page 6

INSULIN THERAPY (Continued)

| Student's self-care insulin administration skills: | | | |
|--|--|--|--|
| Yes No Independently calculates and g | | | |
| | | | |
| Yes No Requires school nurse or traine injections | ed diabetes personnel to calculate/give | | |
| ADDITIONAL INFORMATION FOR STUDEN | T WITH INSULIN PUMP | | |
| Brand/Model of pump: Type | e of insulin in pump: | | |
| Basal rates during school: | | | |
| Type of infusion set: | | | |
| For blood glucose greater than mg/dI hours after correction, consider pump parents/guardian. | that has not decreased within failure or infusion site failure. Notify | | |
| For infusion site failure: Insert new infusion se | et and/or replace reservoir. | | |
| For suspected pump failure: suspend or remov pen. | e pump and give insulin by syringe or | | |
| Physical Activity | | | |
| May disconnect from pump for sports activities | Yes No | | |
| Set a temporary basal rate Yes No Suspend pump use Yes No | | | |
| Student's self-care pump skills: | Independent? | | |
| Count carbohydrates | Yes No | | |
| Bolus correct amount for carbohydrates consumed | Yes No | | |
| Calculate and administer correction bolus | Yes No | | |
| Calculate and set basal profiles | Yes No | | |
| Calculate and set temporary basal rate | Yes No | | |
| Change batteries | Yes No | | |
| Disconnect pump | Yes No | | |
| Reconnect pump to infusion set | Yes No | | |
| Prepare reservoir and tubing | Yes No | | |
| Insert infusion set | Yes No | | |
| Troubleshoot alarms and malfunctions | Yes No | | |

| Diabetes Medical Manag | ement Plan (DMMI | P) – page 7 | |
|--|---------------------------------------|-----------------------|----------------|
| OTHER DIABETES M | EDICATIONS | | |
| Name: | Dose: | Route: _ | Times given: |
| Name: | | | |
| MEAL PLAN | | | |
| Meal/Snack | Time | Carbohydrate Conten | t (grams) |
| Breakfast | | to | |
| Mid-morning snack | · · · · · · · · · · · · · · · · · · · | to | |
| | | to | |
| Mid-afternoon snack | | | |
| Other times to give snack | s and content/amo | ount: | |
| Instructions for when foo sampling event): | - | | 2 0 |
| Special event/party food | | | |
| | | adent discretion | |
| Student's self-care nutrit | _ | | |
| | | arhahydratas | |
| Yes No Independently counts carbohydrates | | | |
| Yes No May count carbohydrates with supervision | | | |
| Yes No Requires school nurse/trained diabetes personnel to count carbohydrates | | | |
| PHYSICAL ACTIVITY | AND SPORTS | | |
| A quick-acting source of juice must be available at | | | |
| Student should eat 1 | 5 grams 🔲 30 g | grams of carbohydrate | other |
| before every 30 |) minutes during | after vigorous phy | sical activity |
| other | | | |

blood ketones are moderate to large. (Additional information for student on insulin pump is in the insulin section on page 6.)

Avoid physical activity when blood glucose is greater than _____ mg/dL or if urine/

If most recent blood glucose is less than ____ mg/dL, student can participate in physical activity when blood glucose is corrected and above ____ mg/dL.

Diabetes Medical Management Plan (DMMP) - page 8

DISASTER PLAN

| To prepare for an unplanned disaster or emergency (72 HOURS), obtain emergency supply kit from parent/guardian. | | | | | |
|---|---|--|--|--|--|
| Continue to follow orders contained in t | his DMMP. | | | | |
| | | | | | |
| Additional insulin orders as follows: Other: | | | | | |
| | | | | | |
| SIGNATURES | | | | | |
| This Diabetes Medical Management Plan ha | s been approved by: | | | | |
| | | | | | |
| Student's Physician/Health Care Provider | Date | | | | |
| I, (parent/guardian:) | give permission to the school nurse | | | | |
| or another qualified health care professional | | | | | |
| (school:) | to perform and carry out the diabetes care | | | | |
| tasks as outlined in (student:) | | | | | |
| Plan. I also consent to the release of the info | | | | | |
| Management Plan to all school staff member | rs and other adults who have responsibility | | | | |
| for my child and who may need to know this | s information to maintain my child's health | | | | |
| and safety. I also give permission to the scho | ool nurse or another qualified health care | | | | |
| professional to contact my child's physician/health care provider. | | | | | |
| | | | | | |
| Acknowledged and received by: | | | | | |
| | | | | | |
| Student's Parent/Guardian | Date | | | | |
| Student's Parent/Guardian | Date | | | | |
| School Nurse/Other Qualified Health Care F | Personnel Date | | | | |